

**JAST (Journal of Animal Science and Technology) TITLE PAGE**  
**Upload this completed form to website with submission**

<b>ARTICLE INFORMATION</b>	<b>Fill in information in each box below</b>
<b>Article Type</b>	Research article
<b>Article Title (within 20 words without abbreviations)</b>	Effects of photoperiod and light intensity on milk production and milk composition of dairy cows in automatic milking system
<b>Running Title (within 10 words)</b>	Light effects on the milk production in AMS
<b>Author</b>	Dong-Hyun Lim <sup>1</sup> , Tae-Il Kim <sup>1</sup> , Sung-Min Park <sup>1</sup> , Kwang-Seok Ki <sup>1</sup> and Younghoon Kim <sup>2</sup>
<b>Affiliation</b>	1 Dairy Science Division, National Institute of Animal Science, Rural Development Administration, Chungcheongnam-do, Cheonan 31000, Korea 2 Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea
<b>ORCID (for more information, please visit <a href="https://orcid.org">https://orcid.org</a>)</b>	Dong-Hyun Lim(0000-0002-8575-0198) Tae-Il Kim(0000-0003-0532-6010) Sung-Min Park(0000-0001-9534-0216) Kwang-Seok Ki(0000-0003-0971-1389) Younghoon Kim (0000-0001-6769-0657)
<b>Competing interests</b>	No potential conflict of interest relevant to this article was reported.
<b>Funding sources</b> State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This study was carried out with the support of “Cooperative Research Program for Agricultural Science and Technology Development (Project title: Development of feeding management technology to increase the melatonin concentration of milk in dairy cattle; Project No: PJ01252001) of the Rural Development Administration of the Republic of Korea).
<b>Acknowledgements</b>	Not applicable.
<b>Availability of data and material</b>	Upon reasonable request, the datasets of this study can be available from the corresponding author.
<b>Authors' contributions</b> Please specify the authors' role using this form.	Conceptualization: Lim DH, Kim TI, Park SM, Ki KS, Kim Y. Data curation: Lim DH, Kim TI, Park SM, Ki KS, Kim Y. Formal analysis: Lim DH, Kim TI, Park SM, Ki KS, Kim Y. Methodology: Lim DH, Kim TI, Park SM, Ki KS. Software: Lim DH, Kim TI, Park SM, Ki KS. Validation: Lim DH, Kim TI, Park SM, Ki KS, Kim Y. Investigation: Lim DH, Kim TI, Park SM, Ki KS, Kim Y. Writing - original draft: Lim DH, Kim TI, Park SM, Ki KS, Kim Y. Writing - review & editing: Lim DH, Kim TI, Park SM, Ki KS, Kim Y.
<b>Ethics approval and consent to participate</b>	All dairy cows were maintained as stated in standard guideline, and the experimental protocol involved in this experiment was approved by the Institutional Animal Care and Use Committee (IACUC) at NIAS (study approval number: IACUC 2017-252).

5 **CORRESPONDING AUTHOR CONTACT INFORMATION**

<b>For the corresponding author (responsible for correspondence, proofreading, and reprints)</b>	<b>Fill in information in each box below</b>
First name, middle initial, last name	Younghoon Kim

Email address – this is where your proofs will be sent	ykeys2584@snu.ac.kr
Secondary Email address	
Address	Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea.
Cell phone number	+82-10-4135-2584
Office phone number	+82-2-880-4808
Fax number	+82-2-873-2271

ACCEPTED

**Running title:** Light effects on the milk production in AMS

10      **Effects of photoperiod and light intensity on milk production and milk  
composition of dairy cows in automatic milking system**

Dong-Hyun Lim<sup>1</sup>, Tae-Il Kim<sup>1</sup>, Sung-Min Park<sup>1</sup>, Kwang-Seok Ki<sup>1</sup> and Younghoon Kim<sup>2,\*</sup>

15      <sup>1</sup>*Dairy Science Division, National Institute of Animal Science, Rural Development Administration,  
Chungcheongnam-do, Cheonan 31000, Korea*

<sup>2</sup>*Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science,  
Seoul National University, Seoul 08826, Korea*

20      \*To whom correspondence should be addressed: [ykeys2584@snu.ac.kr](mailto:ykeys2584@snu.ac.kr)

## ABSTRACT

25

The purpose of this study was to determine the effects of photoperiod and light intensity on milk production, milk composition, hormones levels and blood metabolites indices of Korean Holstein dairy cows in automatic milking system (AMS). A total of 24 Holstein dairy cows were selected and used to four subsequent treatments for the experimental periods of 60 days. The light programs consisted of (1) Control: the natural photoperiod with 14.2 h of the light period and 9.4 h of the dark period (below 10 Lux); (2) T1: 16 hrs of the long day photoperiod (LDPP) with 50 Lux of light; (3) T2: 16 h of LDPP with 100 Lux of light; and (4) T3: 16 h of LDPP with 200 Lux of light, respectively. Importantly, there was a significant difference in the thurl activity of dairy cows between the different light intensity programs ( $p < 0.05$ ). Milk yield was higher in T1 and T2 ( $40.80 \pm 1.71$  and  $39.90 \pm 2.02$  kg/d, respectively) than those of Control and T3 ( $32.18 \pm 1.51$  and  $35.76 \pm 2.80$  kg/d, respectively) ( $p < 0.05$ ), but DMI was lower in T1, T2, and T3 compared to Control ( $p < 0.05$ ). Also, milk fat percentage and the contents of milk fat and total solids were higher in T2 than those in the others ( $p < 0.05$ ). The average daily melatonin level in milk was high to T3 ( $28.20 \pm 0.43$  pg/ml), T2 ( $24.62 \pm 0.32$  pg/ml), T1 ( $19.78 \pm 0.35$  pg/ml), and Control ( $19.36 \pm 0.45$  pg/ml) in order ( $p < 0.05$ ). Also, the cortisol levels in milk and blood were lower in treatment groups than in Control ( $p < 0.05$ ). The results of this study showed that it will be effective to improve the milk yield and milk composition, and to reduce the stress of dairy cows when the light conditions regulate to extend the photoperiod to 16 h at a light emitting diode (LED) intensity of 100 Lux under the AMS in dairy farm.

30

35

40

**Keywords:** Photoperiod; Light intensity; Automatic Milking System; Milk production, Melatonin

45

## INTRODUCTION

Photoperiod is the time period of daily exposure that an organism receives from daylight or artificial light. The photoperiod length has a clear physiological response to reproduction, growth, lactation and health [1]. The intensity of illumination is also known to affect both behavior and physiology of cows [2-5]. Some studies reported that an extended photoperiod could result in an increased milk yield compared to a short photoperiod [6, 7]. Cows exposed to long photoperiod have an increased milk production by 5% to 15% compared to cows held in short photoperiod [6, 8-10]. However, there was no effect on milk yield of dairy cows exposed to lighting for 24-h compared to a natural photoperiod [11]. Phillips and Schofield [12] reported that cows exposed to 481 Lux increased dry matter intake (DMI), milk yield and the time of social activities, while cows exposed to the natural light intensity reduced the lying time. The automatic milking system (AMS) was first introduced into Korea in 2006 [13]. The AMS is in use for 24 h and are based on voluntary visits to the milking unit several times a day [14]. It has the advantage of freeing dairy farmers from labor and time constraints, a greater milk yield and the ability to collect various information on lactating dairy cattle, compared to conventional milking system (CMS) [13, 14]. In contrast to these advantages, previous studies have been conducted on stress in dairy cows milked in barns with an AMS [15-18]. To facilitate cows' visits to the AMS throughout the night, most dairy farmers provide artificial lighting in the waiting area in front of the AMS and in the AMS unit [19]. The number of milking increases when sufficient illumination is maintained compared to guiding light in the barn at night [3], and then a more frequent milking can enhance milk production [20]. On the other side, the exposure to continuous light in the dark period does not have an effect on milk production of dairy cows [11].

Recently, dairy farmers are seeking the management strategy using lighting tool because this approach might be more safe, non-invasive, and effective method to increase milk yield [21-23]. However, what light period and light intensity that is suitable to maximize the milk yield in Korean dairy farms with AMS is not yet fully studied. The proper light conditions for dairy cows, which are

housed in AMS, are very important, as showed many studies that light affects the physiology and behavior of cows. Therefore, this study was performed to evaluate the effects of photoperiod and light  
75 intensity on milk production and composition. Also, it was investigated to determine the variation on the activity, hormone and biochemical indices of dairy cows during different light conditions.

ACCEPTED

## MATERIALS AND METHODS

### 80 **Animals, experimental design and feeding management**

The experiment was carried out at Department of Animal Resources Development, National Institute of Animal Science (NIAS; Cheonan, Republic of Korea). All dairy cows were maintained as stated in standard guideline, and the experimental protocol involved in this experiment was approved by the Institutional Animal Care and Use Committee (IACUC) at NIAS (study approval number: IACUC  
85 2017-252). A total of 24 multiparous Holstein lactating dairy cows (mean  $\pm$  SD,  $2.4 \pm 0.34$  parity) were selected with the average days in milk (DIM)  $114 \pm 44$  DIM, the 7-d milk yield before starting the study was  $35.20 \pm 1.76$  kg/d, the average body weight (BW)  $738.2 \pm 19.7$  kg, and experiment carried out from May to June 2018. Cows were subjected to the same management procedures and housed in a loose  
90 barn. The barn was designed with AMS (Astronaut A3, Lely Industries N.V., Maassluis, the Netherlands) and was modified to control light exposure. All cows had permission to enter the AMS every 4 h or if cows visiting to AMS within 4 h after milking were directly sent to the barn area without letting her stay in the AMS.

Dairy cows were allocated to four subsequent experimental treatments. The treatments were different to the light conditions; (1) Control: the natural photoperiod, which was average 14.2 h of the light period  
95 and 9.4 h of the dark period (below 10 Lux under natural conditions); (2) T1: the long day photoperiod (LDPP) was extended to light with intensity of 50 Lux ( $48.5 \pm 4.7$  Lux); (3) T2: with intensity of 100 Lux ( $104.4 \pm 6.7$  Lux); (4) T3: with intensity of 200 Lux ( $202.8 \pm 9.4$  Lux), respectively. T1, T2, and T3 groups extended the photoperiod by turning on the light emitting diode (LED) at from 04:30 to 07:00  
100 h and 18:00 to 20:30 h. Cows for each treatment group control were exposed to natural light without artificial light or LED light. Cows for LDPP treatment were acclimated to particular light intensity types for 10 days before initiating each treatment period of 5 days. Milk and blood samples were collected at each milking during the period of each experimental treatment.

The nutrient content of the concentrate and TMR samples were analyzed by Foundation of Agri.

Tech. Commercialization & Transfer (Iksan, Republic of Korea). All samples were analyzed by AOAC  
105 [24] for concentrations of moisture, crude fiber, ether extract, crude fiber and crude ash, and by Van  
Soest et al. [25] for concentrations of neutral detergent fiber (NDF) and acid detergent fiber (ADF).  
Total digestible nutrients (TDN), and NEL were calculated with the equations proposed by the NRC  
[26].

All dairy cows fed the same total dietary nutrient provision when considering the sum (NEL1.7  
110 Mcal/kg, and TDN 68.7%) of the total mixed rations and the AMS concentrate. TMR was offered once  
a day at 09:00 h for ad libitum intake, and were fed concentrates according to the milk yield of each  
cow in the special feeder when were milked. The dry matter intake (DMI, kg/day) was estimated at the  
herd level for each group daily as the difference between the amount of feed intake and feed refusal.  
The chemical composition of the rations based on the realized TMR and concentrate are presented in  
115 Table 1.

#### **Measurement of Temperature humidity index (THI)**

The measurement of ambient temperature ( $^{\circ}\text{C}$ ) and relative humidity (RH, %) was monitored with a  
thermo-hygrometer (Testo, model 174H, West Chester, PA) with an accuracy of  $\pm 0.5^{\circ}\text{C}$ , and  $\pm 3\%$  RH.  
120 The thermo-hygrometer was set to record every day per 30 min and placed about 2 meters apart from  
the feeding area. The temperature and humidity values were used to calculate several THI values; THI  
was calculated for each 30 min temperature and humidity measurement according to the formula:  $\text{THI}$   
 $= (0.8 \times ^{\circ}\text{C}) + [\text{RH} \% \times (^{\circ}\text{C} - 14.4)] + 46.4$ , according to Zähler [27].

#### **125 Light control and measurement of activity volume**

Lights of T1, T2, and T3 groups were exposed to cows under the long photoperiod treatment (day  
time : night time = 16 : 8 h) by LED lamps (AFL0312-40W-57KCP123B, AIRTEC SYSTEM CO.,  
LTD, Korea), and controlled by an automatic timer (from 04:30 to 07:00 h and from 18:00 to 20:30 h).  
The loose barn (13  $\times$  50 m) with AMS was installed as followed: 4  $\times$  9 lines, 36 LEDs (Fig. 1). The



130 photo-intensity during the night (21:00 h) was measured with a light meter (Testo, model 540, Testo  
AG, Germany) at intervals of two meters in barn and at cow eyes level (90 cm from the floor).  
Fluorescent and metal halide lights are used as common light sources in dairy facilities [28]. However,  
this study was used LED light, since its lifespan is approximately 12 times longer than that of  
fluorescent lights [29]. Recently, the number of farms using LED lights has been increasing as  
135 installation costs are reduced and long lifespan can decrease dangerous work to replace lights in high  
ceiling of barn.

Neck and thurl activity were measured daily in individual cows with method described by Lim et al.  
[30]. These activity volumes (unit) were calculated using the pedometer (YAMASA Corporation,  
Tokyo, Japan) attached to neck and thurl part of a cow from 09:00 to 18:00 h.

140

#### **Sampling and analysis of blood and milk**

The blood sample was collected via jugular venipuncture of each cow at 14:00 h once a week using  
sterile vacutainer tubes (BD Vacutainer, BD, USA). The collected blood was centrifuged at 1,000 x g  
for 15 min at 4°C. The collected serum was stored at -20°C until analysis. Serum samples were used to  
145 analyze the metabolic indices status (Wako Chemicals, Neuss, Germany) by using blood auto-analyzer  
(Hitachi, 7180, Japan).

During the experimental period, the milk yield was recorded every day with AMS, and milk samples  
of each cow collected for 24 h. The collected milk was used to analyze milk fat, protein, lactose, and  
milk urea nitrogen (MUN) using with LactoScop (MK2, Delta Instruments, the Netherlands). Fat and  
150 protein corrected milk (FPCM) was calculated according to the formula  $FPCM = (0.337 + 0.116 \times \text{Fat } \% + 0.06 \times \text{Protein } \%) \times \text{milk yield (kg/d)}$ . The biochemical indices of glucose, urea nitrogen (UN), Non-  
esterified fatty acids (NEFA), total protein, albumin, and triglycerides (TG) level were analyzed with  
Clinical Analyzer (Hitachi 7180, Japan). The cortisol and melatonin concentration of milk and blood  
were measured using a commercial ELISA kit (Wuhan Abebio Science, Wuhan, China) according to  
155 manufacturer's instructions.

## Statistical Analysis

160 The air temperature, relative humidity, THI, and the duration time of day and night were recorded by date. Also, the activity of neck and leg in dairy cow, the feed intake of TMR and concentrates, body weight, milk yield, and fat, protein, lactose, and MUN of individual milk were recorded. All the raw data were prepared for Microsoft Excel (Microsoft Corp., Redmond, WA, USA) and then analyzed with the statistical package SAS Enterprise Guide 7.1 (SAS Institute Inc., Cary, NC, USA).

165 In order to analyze the differences between light period and light intensity within the same analytical parameters, one-way analysis of variance (ANOVA procedure) was applied for all compositions of Control and treatment groups (included 3 groups exposed to 50, 100, and 200 Lux). Also, the effects of milking time and light intensity on melatonin and cortisol concentration of milk were analyzed with multiple analysis of variance (GLM procedure). Statistical relationships were regarded as being significant when the p value was  $< 0.05$ . A multiple comparison test (Tukey Pairwise Comparisons) was performed to differentiate the mean values of treatments when found significant.

170

## RESULTS AND DISCUSSION

### Weather and photoperiod conditions

175 Average weather conditions and photoperiod trends during the study period are shown in Fig. 2 and Table 2. The estimated THI values averaged 65.65 per day and increased from May (62.58) to June (68.82) ( $p < 0.05$ ). The average daily air temperature ( $T_a$ , °C) and relative humidity (RH, %) were 19.68°C and 66.01 %, respectively.  $T_a$  was increased from May (17.52°C) to June (21.92°C) ( $p < 0.05$ ), while RH was lower in June (64.98 %) than in May (67.31 %) ( $p < 0.05$ ). To date, it has well-established that heat stress during lactation negatively affects milk production. Several studies differently defined on the thermal comfort zone for cows, which Armstrong [31] used  $THI < 71$ , and De Rensis et al. [32] used  $THI < 68$ . Compared with the results of these studies, the average THI value of the present study was investigated for environmental conditions that did not cause to heat stress.

180 In the study period, the daily daytime in June increased by an average of 31 min from May, making the night time in June shorter than in May. Also, the average daytime per day was 14:07 in May and 14:38 in June. Many studies reported that dairy cows exposed to long-day photoperiod (LDPP, 16 h of light) have an increased milk production compared to cow exposed to short-day photoperiod (SDPP, 8 h of light) [6, 9, 10]. Based on these results, treatment groups (T1, T2, and T3) in this study were exposed to the photoperiod (day : night = 16 : 8 h), which was 2 h longer than the control of the natural conditions. However, there were concerns about whether the extension of light interfered with the cow's sleeping hours. Cows sleep total 4 h per day [33], and spend more time sleeping at night compared to day time [34]. Therefore, 8 h of night time for this study was considered as a proper time that did not adversely affect the sleeping time of cows.

### Neck and thurl activity

195 The activity volume of neck and thurl part in day time from 09:00 to 18:00 h with the photoperiod and light intensity are shown in Table 3. The mean activity volume of neck during the day time reduced

with increasing light intensity from 50 to 200 Lux compared with the Control ( $p < 0.05$ ), and the thurl activity of T1 (370.8 unit) and T2 (361.8 unit) was higher than in that of Control (161.0 unit) and T3 (118.4 unit) ( $p < 0.05$ ). Adamczyk et al. [35] reported that mean 24-h activity of cows in early and late lactation remained at a similar level, but appeared to slightly higher relationship between milk yield and activity. In this study, the higher thurl activity of T1 and T2 may be associated with an increase in milk yield, as shown by Adamczyk et al. [35]. Phillips and challengers [4] reported that an optimal level of illumination for walking through the passageways in the dark should be between 39 Lux and 119 Lux. Also, Pettersson and Wiktorsson [3] found that there was no significant difference in cattle preference on the lying area where fully lit or lit with guiding lights only during the dark period.

#### **DMI, BW, milk production and milk composition**

The DMI, BW, milk production, and milk composition trends in different light programs are shown in Table 4. Milk yield in this study was higher in longer photoperiod (50, 100, and 200 Lux, 40.80 kg, 39.90 kg, and 35.76 kg per day, respectively) than in natural photoperiod (Control, 32.18 kg/d) ( $p < 0.05$ ). Then 50 and 100 Lux of light intensity increased compared to 200 Lux group ( $p < 0.05$ ). Among the milk compositions, milk fat percentage was higher in 100 Lux (T2, 4.35%), but was lower in 50 Lux (T1, 3.57%) and 200 Lux (T3, 3.70%) than in Control (3.86%) ( $p < 0.05$ ). The contents of milk fat and total solids were higher at 100 Lux (T2, milk fat 1.62 kg and total solids 4.91 kg per day) than at the others (milk fat 1.15 to 1.30 kg, and total solids 3.94 to 4.57 kg per day) ( $p < 0.05$ ).

Dairy farms use for the daily photoperiod and light intensity system as management tool to improve milk production. Dairy cows exposed to LDPP in lactation period produce more milk yield with 10 to 15% [10] or with 2.5 kg/cow per day [6] compared to cows exposed to SDPP. Similar to previous studies, the present study also appeared to increase the milk yield with 15.9% at 50 Lux and with 13.4% at 100 Lux in longer photoperiod (16 h of light per day). The exposure to continuous light in the dark period does not have a positive effect on milk production of dairy cows [11]. In this regard, Buchanan et al. [36] suggested that a dark period is necessary to maintain the photoperiodic responses, since cows

exposed in continuous lighting may be lost the ability to recognize the day length. Especially during the night intensity of dairy cows, Muthuramalingam et al. [2] reported less than 10 Lux, while Bal et al. [37] suggested 40 to 60 Lux.

The conflicting results were reported by previous studies on the milk composition obtained by the different photoperiod. Miller et al. [38] reported that milk composition was not affected by photoperiod management. Bodurov [8] found that milk fat content increased by 0.3% in LDPP, however, other studies reported milk fat percentage decreased to LDPP [6, 12]. This study found that the milk fat content increased as milk yield increased to the LDPP conditions compared to the natural photoperiod, although it was difficult to identify the effect of the light intensity on milk fat percentage.

Some studies reported that cows exposed to LDPP increased by 0.8 to 1.5 kg/d of DMI to support the higher milk production [9, 38]. Prior to conducting this study, the DMI was expected to be higher as milk yield increased to the treatment groups, but in fact, DMI in the control group was higher. This result was supported by Peters et al. [10] reported that additional light and longer light period (16 h of 114 to 207 Lux vs. 9 or 12 h of 39 to 93 Lux) were increased both growth and milk yield without any increase in feed consumption. This could be explained that the time for intake and conversion of feed and the productivity are influenced at a higher extent by the physiological state and social hierarchy than by the photoperiod [39].

240

#### **Melatonin and cortisol concentrations of milk**

Milk melatonin and cortisol concentrations milked with AMS according to the milking time per 24-h are shown in Table 5. Melatonin is a neuro-hormone derived from serotonin during the dark phase, and produced particularly in the pineal gland, but also in the retina of vertebrates [40]. In this study, average daily melatonin level in milk was higher in treatment groups than in control, and increased to T3 (28.20 pg/ml), T2 (24.62 pg/ml), and T1 (19.78 pg/ml) in order ( $p < 0.05$ ). These results exhibited that the daily melatonin concentration in milk was high in LDPP than in natural photoperiod, and increased as the light intensity increased from 50 to 200 Lux. Milk melatonin level was different with

the milking times that it was high at 08:01 to 12:00 h in Control, at 16:01 to 20:00 h in T1 and T2, at  
250 20:01 to 24:00 h in T3 ( $p < 0.05$ ). Vanecek [41] reported that the duration of melatonin increase was  
short on long day photoperiods and long on short day photoperiods. These results were different with  
the current study that the melatonin level in milk retained longer in T2 and T3 compared to milk  
melatonin level at 08:01 to 12:00 h in Control. Moreover, cortisol which is a hormone of glucocorticoid  
class, is sensitively responded to light with a distinct circadian rhythm, and is one of the most important  
255 stress indicators in mammals [42]. Cortisol concentration in milk was higher in control than that in  
treatment groups (T1, T2, and T3) ( $p < 0.05$ ). Average cortisol concentration was lowest in T1 exposed  
to 50 Lux among the treatment groups under the LDPP condition of daytime from 04:30 to 20:30 h ( $p$   
< 0.05).

#### 260 **Biochemical indices in blood**

The biochemical indices level in blood of dairy cows exposed under the different light programs is  
shown in Table 6. The level of blood melatonin was lower in T2 (17.44 pg/ml) and T3 (17.03 pg/ml)  
than in Control (25.55 pg/ml) and T1 (23.72 pg/ml) ( $p < 0.05$ ). Melatonin is a relationship between  
blood and milk concentrations [41], since melatonin is amphiphilic, so it can freely diffuse through  
265 biological membranes into the circulatory system and from the bloodstream into the milk [43].  
Kollmann et al. [44] found about 40% of the blood melatonin concentration in the milk of cows  
producing approximately 32 kg/milk/day. However, this study showed a different tendency to melatonin  
level in blood collected at 14:00 h compared to milk milked at 12:01 to 16:00 h. Blood cortisol  
concentration decreased the treatment groups compared with the Control ( $p < 0.05$ ). These results could  
270 be explained by previous studies [45, 46]. Exposed to light stimulates the gene expression in adrenal  
gland causing plasma corticosterone surge in mammals Ishida et al. [45]. Hyder et al. [46] suggested  
that melatonin may inhibit this gene expression to reduce cortisol secretion. Taken together, we consider  
that dairy cows may be produced more melatonin in body fluids under the LDPP than that under the  
SDPP, and then could relieve their stress.

275 Blood metabolites can indicate the energy and protein metabolism and liver health of the dairy cows  
[47]. In this study, BUN level in blood was significantly increased for lactating dairy cows exposed to  
LDPP compared to those exposed to natural photoperiod. Creatinine level was showed the similar  
tendency to BUN, which is positively correlated with MUN and creatinine level [48]. Previous studies  
found that some blood metabolites were variable for animals exposed to different light photoperiod and  
280 color [28, 49]. On that reason, these studies suggested because cows exposed to long photoperiod might  
be attributed to greater feed intake. However, DMI in this study decreased to dairy cows exposed to  
LDPP than those exposed to natural conditions. Previous studies tried to explain the reason that milk  
yield and blood metabolites were higher for LDPP without increasing DMI [10, 28, 39]. Espinoza [28]  
suggested that the daily rhythms of blood metabolism may be altered by the circadian rhythms such as  
285 the light-dark cycle.

Dahl [50] suggested that the milk production of dairy cows increased at the height of about 91 cm  
above the stall floor at 150 Lux of light intensity, and also increased in LDPP, which is 16 to 18 h of  
light followed by 6 to 8 h of darkness in a 24-h period [6]. The results of this study showed that it will  
be effective in reducing stress on dairy cows and improving milk productivity and milk compositions  
290 (fat and protein) when the light conditions regulate to extend the photoperiod to 16 h at a LED intensity  
of 100 Lux in dairy farm, which has a AMS, compared to natural light conditions. The difference in the  
present study compared with previous studies may be due to the geographical and environmental  
differences. Latitude affects the incidence angle of solar radiation and the length of photoperiod.  
Daylight period is longest at the summer solstice and shortest at the winter solstice in the Northern  
295 Hemisphere [51]. Therefore, further research is needed to optimize the light conditions in order to  
improve the milk productivity of dairy cows housed in loose barn with AMS in consideration of Korea's  
geographical environment

## CONCLUSIONS

300 This study discussed effects of light intensity and light period on yield and compositions of milk,

stress related hormones and biochemical indices in Holstein dairy cows, which milked in AMS. The 50 and 100 Lux exposed dairy cows showed more milk yield than those of other groups. Also, the 100 Lux exposed dairy cows exhibited higher level of fat, protein and total solids in milk as compared to other group cows. The level of melatonin in milk was significantly increased as the light intensity increased.

305 Whereas, the cortisol levels in milk was lower in treatment groups than in Control. Our results suggest that the difference of photoperiod and light intensity could act as external stimulation to the rhythmic pattern (metabolites) involved in alteration of hormones function, milk yield, and milk compositions. Additionally, these results of the study are considering the widespread use of photoperiod in dairy animal industry to increasing incidence of antioxidant levels. In recent, it has been reported that the  
310 regulation of circadian rhythms via photoperiod and light intensity significantly influences on the performance and physiology of dairy cows [52]. Ongoing study is evaluating on molecular mechanism and circadian pattern underlying how light intensity and light period affect the milk production and compositions.

### 315 **Acknowledgements**

This study was carried out with the support of “Cooperative Research Program for Agricultural Science and Technology Development (Project title: Development of feeding management technology to increase the melatonin concentration of milk in dairy cattle; Project No: PJ01252001) of the Rural Development Administration of the Republic of Korea).

320



## References

1. Dahl G, Tao S, Thompson I. Lactation Biology Symposium: Effects of photoperiod on mammary gland development and lactation. *Journal of animal science*. 2012;90(3):755-60.
2. Muthuramalingam P, Kennedy A, Berry R. Plasma melatonin and insulin-like growth factor-1 responses to dim light at night in dairy heifers. *Journal of pineal research*. 2006;40(3):225-9.
3. Pettersson D, Wiktorsson H. Illumination or guiding light during night hours in the resting areas of AM-barns. *Automatic Milking-a better understanding*, Meijering, A, Hogeveen, H, de Koning CJAM,(Eds). 2004:468-73.
4. Phillips C, Morris I, Lomas C, Lockwood S. The locomotion of dairy cows in passageways with different light intensities. *Animal welfare*. 2000;9(4):421-31.
5. Stanisiewski E, Chapin L, Ames N, Zinn S, Tucker H. Melatonin and prolactin concentrations in blood of cattle exposed to 8, 16 or 24 hours of daily light. *Journal of animal science*. 1988;66(3):727-34.
6. Dahl G, Buchanan B, Tucker H. Photoperiodic Effects on Dairy Cattle: A Review<sup>1</sup>. *Journal of dairy science*. 2000;83(4):885-93.
7. Dahl G, Petitclerc D. Management of photoperiod in the dairy herd for improved production and health. *Journal of animal science*. 2003;81(15\_suppl\_3):11-7.
8. Bodurov N. Effect of supplementary artificial illumination with visible rays on biochemical indices in the blood serum, milk yields and fertilization during lactation. *Veterinarno-Meditsinski Nauki*. 1979;16(6):58-65.
9. Collier RJ, Dahl G, VanBaale M. Major advances associated with environmental effects on dairy cattle. *Journal of dairy science*. 2006;89(4):1244-53.
10. Peters R, Chapin L, Leining K, Tucker H. Supplemental lighting stimulates growth and lactation in cattle. *Science*. 1978;199(4331):911-2.
11. Marcek J, Swanson L. Effect of photoperiod on milk production and prolactin of Holstein dairy cows. *Journal of Dairy Science*. 1984;67(10):2380-8.
12. Phillips C, Schofield S. The effect of supplementary light on the production and behaviour of dairy cows. *Animal Science*. 1989;48(2):293-303.
13. Lee J-S, Nam K-T, Park S-M, Son Y-S. Questionnaire study on the difficulties and improvements of the 6th Industrialization Dairy Farm. *Journal of Dairy Science and Biotechnology*. 2016;34(4):255-62.
14. Jacobs J, Siegford J. Invited review: The impact of automatic milking systems on dairy cow management, behavior, health, and welfare. *Journal of dairy science*. 2012;95(5):2227-47.
15. Gygax L, Neuffer I, Kaufmann C, Hauser R, Wechsler B. Restlessness behaviour, heart rate and heart-rate variability of dairy cows milked in two types of automatic milking systems and auto-tandem milking parlours. *Applied Animal Behaviour Science*. 2008;109(2-4):167-79.
16. Hagen K, Lexer D, Palme R, Troxler J, Waiblinger S. Milking of Brown Swiss and Austrian Simmental cows in a herringbone parlour or an automatic milking unit. *Applied Animal Behaviour Science*. 2004;88(3-4):209-25.
17. Hopster H, Bruckmaier R, Van der Werf J, Korte S, Macuhova J, Korte-Bouws G, et al. Stress responses during milking; comparing conventional and automatic milking in primiparous dairy cows. *Journal of Dairy Science*. 2002;85(12):3206-16.
18. Wenzel C, Schönreiter-Fischer S, Unshelm J. Studies on step-kick behavior and stress of cows during milking in an automatic milking system. *Livestock Production Science*. 2003;83(2-3):237-46.
19. Helmreich S, Wechsler B, Hauser R, Gygax L. Effects of milking frequency in automatic milking systems on salivary cortisol, immunoglobulin A, somatic cell count and melatonin. *Schweizer Archiv fur Tierheilkunde*. 2016;158(3):179-86.
20. Wright J, Wall E, McFadden T. Effects of increased milking frequency during early lactation on milk yield and udder health of primiparous Holstein heifers. *Journal of animal science*. 2013;91(1):195-202.
21. Auchtung T, Salak-Johnson J, Morin D, Mallard C, Dahl G. Effects of photoperiod during the dry period on cellular immune function of dairy cows. *Journal of dairy science*. 2004;87(11):3683-9.
22. Thomas AA, Thelen JT, Go AS, Surbrook T, Vanece MA, Althouse J. A Dairy Long Day Lighting Success Story: MI Dairy Increases Production and Cuts Costs. 2017 ASABE Annual International Meeting; St. Joseph, MI: ASABE; 2017. p. 1.
23. VanZweden BJ, Thomas AA, Go AS, Surbrook TC. Expanded LED benefits through an automated long day lighting system at a 3x milking dairy farm. 2019 ASABE Annual International Meeting; St.

- 375 Joseph, MI: ASABE; 2019. p. 1.
24. Cunniff P. Official methods of analysis. Association of Official Analytical Chemists (AOAC) 16th ed  
Arlington, Virginia, USA. 1995.
25. Van Soest Pv, Robertson J, Lewis B. Methods for dietary fiber, neutral detergent fiber, and nonstarch  
polysaccharides in relation to animal nutrition. *Journal of dairy science*. 1991;74(10):3583-97.
- 380 26. NRC I. Nutrient requirements of dairy cattle. National Research Council. 2001.
27. Zähler M, Schrader L, Hauser R, Keck M, Langhans W, Wechsler B. The influence of climatic  
conditions on physiological and behavioural parameters in dairy cows kept in open stables. *Animal  
Science*. 2004;78(1):139-47.
28. Espinoza OS, Oba M. Interaction effect of photoperiod management and dietary grain allocation on  
productivity of lactating dairy cows. *Canadian journal of animal science*. 2017;97(3):517-25.
- 385 29. Yeh N, Chung J-P. High-brightness LEDs—Energy efficient lighting sources and their potential in  
indoor plant cultivation. *Renewable and Sustainable Energy Reviews*. 2009;13(8):2175-80.
30. Lim DH, Kim TI, Kim HJ, Kim SB, Park SM, Park JH, et al. Effect of Short-distance Walking Activity  
on Milk Production and Metabolic Status of Lactating Dairy Cows. *Journal of The Korean Society of  
Grassland and Forage Science*. 2018;38(4):343-8.
- 390 31. Armstrong D. Heat stress interaction with shade and cooling. *Journal of dairy science*. 1994;77(7):2044-  
50.
32. De Rensis F, Garcia-Ispuerto I, López-Gatius F. Seasonal heat stress: Clinical implications and hormone  
treatments for the fertility of dairy cows. *Theriogenology*. 2015;84(5):659-66.
- 395 33. Ruckebusch Y. The relevance of drowsiness in the circadian cycle of farm animals. *Animal behaviour*.  
1972;20(4):637-43.
34. Nilsson E. Quantification of sleep in dairy cows in three different stages of lactation. 2011.
35. Adamczyk K, Gil Z, Felenczak A, Skrzyński G, Zapletal P, Choroszy Z. Relationship between milk  
yield of cows and their 24-hour walking activity. *Animal Science Papers and Reports*. 2011;29(3):185-  
95.
- 400 36. Buchanan BA, Chapin LT, Tucker HA. Prolonged suppression of serum concentrations of melatonin in  
prepubertal heifers. *Journal of pineal research*. 1992;12(4):181-9.
37. Bal MA, Penner GB, Oba M, Kennedy AD. Effects of dim light at night on milk yield, milk composition  
and endocrine profile of lactating dairy cows. *Canadian journal of animal science*. 2008;88(4):609-12.
- 405 38. Miller A, Stanisiewski E, Erdman R, Douglass L, Dahl G. Effects of long daily photoperiod and bovine  
somatotropin (Trobect®) on milk yield in cows. *Journal of dairy science*. 1999;82(8):1716-22.
39. Penev T, Radev V, Slavov T, Kirov V, Dimov D, Atanassov A, et al. Effect of lighting on the growth,  
development, behaviour, production and reproduction traits in dairy cows. *Int J Curr Microbiol App Sci*.  
2014;3(11):798-810.
- 410 40. Tosini G, Fukuhara C. Photic and circadian regulation of retinal melatonin in mammals. *Journal of  
neuroendocrinology*. 2003;15(4):364-9.
41. Vanecek J. Cellular mechanisms of melatonin action. *Physiological reviews*. 1998;78(3):687-721.
42. Carlson LE, Campbell TS, Garland SN, Grossman P. Associations among salivary cortisol, melatonin,  
catecholamines, sleep quality and stress in women with breast cancer and healthy controls. *Journal of  
Behavioral Medicine*. 2007;30(1):45-58.
- 415 43. Romanini EB, Volpato AM, Sifuentes dos Santos J, de Santana EHW, de Souza CHB, Ludovico A.  
Melatonin concentration in cow's milk and sources of its variation. *Journal of Applied Animal Research*.  
2019;47(1):140-5.
44. Kollmann M, Locher M, Hirche F, Eder K, Meyer H, Bruckmaier R. Effects of tryptophan  
supplementation on plasma tryptophan and related hormone levels in heifers and dairy cows. *Domestic  
animal endocrinology*. 2008;34(1):14-24.
- 420 45. Ishida A, Mutoh T, Ueyama T, Bando H, Masubuchi S, Nakahara D, et al. Light activates the adrenal  
gland: timing of gene expression and glucocorticoid release. *Cell metabolism*. 2005;2(5):297-307.
46. Hyder I, Sejian V, Bhatta R, Gaughan J. Biological role of melatonin during summer season related heat  
stress in livestock. *Biological Rhythm Research*. 2017;48(2):297-314.
- 425 47. Cozzi G, Ravarotto L, Gottardo F, Stefani A, Contiero B, Moro L, et al. Reference values for blood  
parameters in Holstein dairy cows: Effects of parity, stage of lactation, and season of production.

- Journal of Dairy Science. 2011;94(8):3895-901.
- 430 48. Nozad S, Ramin A-G, Moghadam G, Asri-Rezaei S, Babapour A, Ramin S, editors. Relationship between blood urea, protein, creatinine, triglycerides and macro-mineral concentrations with the quality and quantity of milk in dairy Holstein cows. Veterinary Research Forum; 2012: Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.
- 435 49. Son J, Park J, Kang D, Belal SA, Cha J, Shim K. Effects of white, yellow, and blue colored LEDs on milk production, milk composition, and physiological responses in dairy cattle. Animal Science Journal. 2020;91(1):e13337.
50. Dahl GE. Let there be light: Photoperiod management of cows for production and health. 2005.
51. Webster J, Corson I, Littlejohn R, Stuart S, Suttie J. Photoperiodic requirements for rapid growth in young male red deer. Animal science. 1998;67(2):363.
- 440 52. Shehab-El-Deen MAM, Fadel MS, Van Soom A, Saleh SY, Maes D, Leroy JL. Circadian rhythm of metabolic changes associated with summer heat stress in high-producing dairy cattle. Tropical Animal Health and Production. 2010;42(6):1119-25.

ACCEPTED

**Table 1.** Chemical composition of the concentrates and TMR fed for dairy cows throughout the study<sup>1</sup>

Items	Concentrates	TMR
Ingredients ratio (% , DM)		
Concentrate	-	17.75
Cashew nut meal	-	8.39
Soybean meal	-	4.15
Corn silage	-	29.16
Mixed hay	-	16.27
Alfalfa	-	12.71
Timothy	-	10.58
Bypass fat	-	0.94
Sodium bicarbonate	-	0.24
Yeast culture	-	0.24
Mineral mixture	-	0.24
Calcium carbonate	-	0.33
Nutrient compositions (% , DM)		
Moisture (% , <i>as fed</i> )	10.92	42.89
Crude protein	18.96	10.72
Ether extract	4.16	3.53
Crude fiber	4.55	14.49
Crude ash	7.93	5.77
Neutral detergent fiber (NDF)	19.01	27.88
Acid detergent fiber (ADF)	7.35	16.14
Calcium	1.40	0.61
Phosphorous	0.57	0.25
NFE (%)	53.48	22.60
TDN (%)	74.18	67.70
NE <sub>L</sub> (Mcal/kg)	1.70	1.54

445 <sup>1</sup>Values obtained from Foundation of Agri. Tech. Commercialization & Transfer (Iksan, Korea).

**Table 2.** The daily average of air temperature (Ta, °C), relative humidity (RH, %), Temperature-humidity index (THI) and photoperiod during the study period

Item	May	June	Mean	SEM	p-value
THI	62.58	68.82	65.65	0.60	<0.001
Ta (°C)	17.52	21.92	19.68	0.41	<0.001
RH (%)	67.31	64.98	66.01	1.45	0.372
Photoperiod (hh:mm)					
Daytime	14:07	14:38	14:22	0.00	<0.001
Nighttime	9:52	9:21	9:37	0.00	<0.001

450

ACCEPTED

**Table 3.** Comparative performance on activity volume<sup>1)</sup> in lactating dairy cows when changing of light intensity

Item	Control	T1	T2	T3	SEM	<i>p</i> -value
Neck part	2904.7 <sup>a</sup>	2171.3 <sup>ab</sup>	1553.3 <sup>bc</sup>	1154.4 <sup>c</sup>	180.7	0.002
Thurl part	161.0 <sup>b</sup>	370.8 <sup>a</sup>	361.8 <sup>a</sup>	118.4 <sup>b</sup>	26.1	0.001

Control, natural photoperiod; T1, LDPP (day : night = 16 : 8 h) with 50 Lux of the light intensity; T2, LDPP with

455 100 Lux; T3, LDPP with 200 Lux.

<sup>1)</sup> Activity volume (unit) was counted using pedometer from 09:00 to 18:00 daily.

<sup>a-c</sup> denotes comparison made within rows ( $p < 0.05$ ).

ACCEPTED

**Table 4.** Dry matter intake, body weight, milk production and milk composition in lactating dairy cows

Item	Control	T1	T2	T3	SEM	p-value
DMI, kg/d	29.03 <sup>a</sup>	27.35 <sup>b</sup>	26.71 <sup>b</sup>	28.90 <sup>ab</sup>	0.26	0.004
Body weight, kg	738.18	733.55	727.27	727.00	9.08	0.968
Milk yield, kg/d	32.18 <sup>b</sup>	40.80 <sup>a</sup>	39.90 <sup>a</sup>	35.76 <sup>ab</sup>	1.03	0.003
FPCM <sup>1)</sup> , kg/d	29.99 <sup>c</sup>	37.29 <sup>ab</sup>	39.57 <sup>a</sup>	33.75 <sup>bc</sup>	0.99	0.003
Milk fat, %	3.86 <sup>b</sup>	3.57 <sup>c</sup>	4.35 <sup>a</sup>	3.70 <sup>bc</sup>	0.04	<0.001
Milk protein, %	3.21 <sup>ab</sup>	3.25 <sup>a</sup>	3.26 <sup>a</sup>	3.16 <sup>b</sup>	0.01	0.066
Lactose, %	4.78 <sup>b</sup>	4.84 <sup>a</sup>	4.79 <sup>b</sup>	4.80 <sup>ab</sup>	0.01	0.033
MUN <sup>2)</sup> , mg/100g	14.97 <sup>c</sup>	19.13 <sup>b</sup>	20.29 <sup>a</sup>	20.21 <sup>a</sup>	0.18	<0.001
Milk fat, kg	1.15 <sup>b</sup>	1.30 <sup>b</sup>	1.62 <sup>a</sup>	1.26 <sup>b</sup>	0.05	0.005
Milk protein, kg	0.99	1.18	1.21	1.08	0.04	0.143
Lactose, kg	1.55	1.81	1.83	1.68	0.06	0.349
Total solids, kg	3.94 <sup>b</sup>	4.57 <sup>ab</sup>	4.91 <sup>a</sup>	4.28 <sup>ab</sup>	0.16	0.137

460 Control, natural photoperiod; T1, LDPP (day : night = 16 : 8 h) with 50 Lux of the light intensity; T2, LDPP with 100 Lux; T3, LDPP with 200 Lux.

<sup>1)</sup> Fat and protein corrected milk (FPCM) was calculated by milk yield  $\times$  (0.337 + 0.116  $\times$  Fat % + 0.06  $\times$  Protein %).

<sup>2)</sup> Milk urea nitrogen.

465 <sup>a-c</sup> denotes comparison made within rows ( $p < 0.05$ ).

**Table 5.** Melatonin and cortisol concentrations in milk of dairy cows milked with automatic milking system according to the milking time per 24-h

Milking time (MT)	Control	T1	T2	T3	SEM	<i>p</i> -value		
						Treat	MT	Treat × MT
..... <i>Melatonin, pg/ml</i> .....								
00:01 ~ 04:00	17.17 <sup>y</sup>	18.82 <sup>yz</sup>	25.63 <sup>xy</sup>	27.58 <sup>xyz</sup>	0.70	<0.001		
04:01 ~ 08:00	19.18 <sup>y</sup>	17.85 <sup>z</sup>	24.73 <sup>y</sup>	28.78 <sup>xy</sup>	0.68	<0.001		
08:01 ~ 12:00	25.64 <sup>x</sup>	20.38 <sup>xy</sup>	21.79 <sup>z</sup>	26.35 <sup>yz</sup>	0.62	<0.001		
12:01 ~ 16:00	19.48 <sup>y</sup>	20.42 <sup>xy</sup>	22.55 <sup>z</sup>	25.22 <sup>z</sup>	0.40	<0.001		
16:01 ~ 20:00	19.28 <sup>y</sup>	22.11 <sup>x</sup>	26.95 <sup>x</sup>	29.60 <sup>wx</sup>	0.71	<0.001		
20:01 ~ 24:00	19.01 <sup>y</sup>	20.51 <sup>xy</sup>	26.00 <sup>xy</sup>	31.86 <sup>w</sup>	0.80	<0.001		
Mean	19.36 <sup>c</sup>	19.78 <sup>c</sup>	24.62 <sup>b</sup>	28.20 <sup>a</sup>	0.28	<0.001	0.007	<0.001
..... <i>Cortisol, pg/ml</i> .....								
00:01 ~ 04:00	1016.25 <sup>y</sup>	777.78 <sup>xy</sup>	906.81 <sup>x</sup>	787.47 <sup>y</sup>	15.51	<0.001		
04:01 ~ 08:00	1041.33 <sup>xy</sup>	798.62 <sup>xy</sup>	886.40 <sup>xyz</sup>	760.71 <sup>y</sup>	14.85	<0.001		
08:01 ~ 12:00	1091.50 <sup>x</sup>	807.69 <sup>xy</sup>	845.80 <sup>yz</sup>	903.67 <sup>x</sup>	20.90	<0.001		
12:01 ~ 16:00	975.07 <sup>y</sup>	764.29 <sup>y</sup>	834.59 <sup>z</sup>	935.34 <sup>x</sup>	12.96	<0.001		
16:01 ~ 20:00	1004.45 <sup>y</sup>	752.09 <sup>y</sup>	889.90 <sup>xy</sup>	884.99 <sup>x</sup>	14.37	<0.001		
20:01 ~ 24:00	998.78 <sup>y</sup>	831.69 <sup>x</sup>	858.52 <sup>xyz</sup>	780.69 <sup>y</sup>	14.71	<0.001		
Mean	1011.14 <sup>a</sup>	790.65 <sup>c</sup>	870.75 <sup>b</sup>	840.63 <sup>bc</sup>	6.29	<0.001	0.049	<0.001

470 Control, natural photoperiod; T1, LDPP (day : night = 16 : 8 h) with 50 Lux of the light intensity; T2, LDPP with 100 Lux; T3, LDPP with 200 Lux.

<sup>a-c</sup> denote comparison made within row

<sup>w-z</sup> denote comparison made within column



475 **Table 6.** Biochemical indices status in lactating dairy cows

Item	Control	T1	T2	T3	SEM	<i>p</i> -value
Melatonin, pg/ml	25.55 <sup>a</sup>	23.72 <sup>a</sup>	17.44 <sup>b</sup>	17.03 <sup>b</sup>	0.78	<0.001
Cortisol, pg/ml	1437.61 <sup>a</sup>	1224.31 <sup>b</sup>	953.03 <sup>c</sup>	935.52 <sup>c</sup>	44.83	<0.001
Total protein, g/dl	55.90	59.20	62.72	61.69	1.45	0.389
Albumin, g/dl	2.28	2.42	2.62	2.63	0.07	0.198
BUN, mg/dl	15.26 <sup>b</sup>	17.94 <sup>a</sup>	18.54 <sup>a</sup>	19.36 <sup>a</sup>	0.40	0.001
Creatinine, mg/dl	1.43	1.46	1.53	1.63	0.03	0.059
Glucose, mg/dl	49.83	49.17	51.92	49.33	0.77	0.590
Non-esterified fatty acids, mg/dl	42.92	29.58	31.92	34.25	3.25	0.531
Triglyceride, mg/dl	4.42	4.75	4.83	5.58	0.23	0.357
G-GTP, IU/L	18.58	24.58	26.83	24.58	2.73	0.768
GOT(AST), IU/L	47.17	52.17	56.25	58.83	2.30	0.331
GPT(ALT), IU/L	14.42 <sup>c</sup>	18.08 <sup>bc</sup>	21.08 <sup>ab</sup>	24.08 <sup>a</sup>	0.83	<0.001
Cholesterol, mg/dl	121.92	137.83	147.17	145.25	5.33	0.352

The blood sample was collected via jugular venipuncture of each cow at 14:00 h.

Control, natural photoperiod; T1, LDPP (day : night = 16 : 8 h) with 50 Lux of the light intensity; T2, LDPP with 100 Lux; T3, LDPP with 200 Lux.

<sup>a-c</sup> denote comparison made within row

480

## FIGURE LEGENDS

**Figure 1.** Schematic layout of barn with LED lights. The photo-intensity during the night (21:00 h) was measured with a light meter at intervals of two meters in barn and at cow eyes level (90 cm from the floor).

485

**Figure 2.** The average daily value of weather conditions including air temperature, relative humidity, and temperature-humidity index (A), and photoperiod trends (B) from May to June.

490

ACCEPTED





Figure 1.

ACCETV



495

Figure 2.

Dear Editor,

We submit the revised paper entitled: “Effects of photoperiod and light intensity on milk production and milk composition of dairy cows in automatic milking system” (jast 2021-00092 Version 1).—

The reviewers’ questions have been underlined and our responses are detailed below.—

#### **REVIEWER 1**

This manuscript describes the photoperiod and light intensity on milk production and milk composition of dairy cows. It is useful information about the insight into dairy cow management. It showed the improving the milk yield, milk fat and total solids in T2 and lower DMI in treatment than control. The stress of dairy cows was reduced by 100 Lux of light. The paper is presenting solid experimental results backed by convincing and in-depth simulations and analysis and I believe that it is a good candidate for being published in Journal on Animal Science and Technology.

—We appreciate on the reviewer’s kind considerations.

#### **REVIEWER 2**

The study investigated the effects of photoperiod and light intensity on milk production, milk composition, hormones levels, and blood metabolites indices of Korean Holstein dairy cows in the automatic milking system.

For a more accurate comparison of the effect of photoperiod, it would be better to add one more treatment, the LDPP with below 10 Lux.

—As suggested, additional treatments (below 10 Lux of LDPP) could be supported to establish on our optimized condition for LDPP. Unfortunately, it’s so hard to control/maintain the intensity of low level of Lux technically. Furthermore, if having next application, we will be employ on the low Lux condition under controlled facilities.

Please insert the symbol of multiply (×) in the caption of Table 4

—As suggested, we carefully revised Table 4.—

Please mention the time when the blood sample is collected in the title of Table 6.

As suggested, we indicated blood sampling time at legend of Table 6.—